

VOLUME 98 PART 5

JULY 1988

ISSN 0303-2515

NK
74 SE
= (52768)

ANNALS

OF THE SOUTH AFRICAN
MUSEUM

CAPE TOWN



INSTRUCTIONS TO AUTHORS

1. MATERIAL should be original and not published elsewhere, in whole or in part.
2. LAYOUT should be as follows:
 - (a) *Centred masthead to consist of*
Title: informative but concise, without abbreviations and not including the names of new genera or species
Author's(s') name(s)
Address(es) of author(s) (institution where work was carried out)
Number of illustrations (figures, enumerated maps and tables, in this order)
 - (b) *Abstract* of not more than 200 words, intelligible to the reader without reference to the text
 - (c) *Table of contents* giving hierarchy of headings and subheadings
 - (d) *Introduction*
 - (e) *Subject-matter* of the paper, divided into sections to correspond with those given in table of contents
 - (f) *Summary*, if paper is lengthy
 - (g) *Acknowledgements*
 - (h) *References*
 - (i) *Abbreviations*, where these are numerous.

3. MANUSCRIPT, to be submitted in triplicate, should be typewritten and neat, double spaced with 3 cm margins all round. First lines of paragraphs should be indented. Tables and a list of captions for illustrations should be typed separately, their positions indicated in the text. All pages should be numbered consecutively.

Major headings of the paper are centred capitals; first subheadings are shouldered small capitals; second subheadings are shouldered italics; third subheadings are indented, shouldered italics. Further subdivisions should be avoided, as also enumeration (never roman numerals) of headings and abbreviations.

Footnotes should be avoided unless they are short and essential.

Only generic and specific names should be underlined to indicate italics; all other marking up should be left to editor and publisher.

4. ILLUSTRATIONS should be reducible to a size not exceeding 12 × 18 cm (19 cm including caption); the reduction or enlargement required should be indicated (and preferably uniform); originals larger than 35 × 47 cm should not be submitted; photographs should be rectangular in shape and final size. A metric scale should appear with all illustrations, otherwise magnification or reduction should be given in the caption; if the latter, then the final reduction or enlargement should be taken into consideration.

All illustrations, whether line drawings or photographs, should be termed figures (plates are not printed; half-tones will appear in their proper place in the text) and numbered in a single series. Items of composite figures should be designated by capital letters; lettering of figures is not set in type and should be in lower-case letters. If Letraset is used authors are requested to use Helvetica-style lettering, if possible.

The number of the figure should be lightly marked in pencil on the back of each illustration.

5. REFERENCES cited in text and synonymies should all be included in the list at the end of the paper, using the Harvard System (*ibid.*, *idem*, *loc. cit.*, *op. cit.* are not acceptable):

- (a) Author's name and year of publication given in text, e.g.:

'Smith (1969) describes . . .'

'Smith (1969: 36, fig. 16) describes . . .'

'As described (Smith 1969a, 1969b; Jones 1971)'

'As described (Haughton & Broom 1927) . . .'

'As described (Haughton *et al.* 1927) . . .'

Note: no comma separating name and year

pagination indicated by colon, not p.

names of joint authors connected by ampersand

et al. in text for more than two joint authors, but names of all authors given in list of references.

- (b) Full references at the end of the paper, arranged alphabetically by names, chronologically within each name, with suffixes *a*, *b*, etc., to the year for more than one paper by the same author in that year, e.g. Smith (1969a, 1969b) and not Smith (1969, 1969a).

For books give title in italics, edition, volume number, place of publication, publisher.

For journal article give title of article, title of journal in italics (according to the *World list of scientific periodicals*, 4th ed. London: Butterworths, 1963), series in parentheses, volume number, part number in parentheses, pagination (first and last pages of article).

Examples (note capitalization and punctuation)

BULLOUGH, W. S. 1960. *Practical invertebrate anatomy*. 2nd ed. London: Macmillan.

FISCHER, P. H. 1948. Données sur la résistance et de la vitalité des mollusques. *Journal de conchyliologie* 88 (3): 100–140.

FISCHER, P. H., DUVAL, M. & RAFFY, A. 1933. Études sur les échanges respiratoires des littorines. *Archives de zoologie expérimentale et générale* 74 (33): 627–634.

KOHN, A. J. 1960a. Ecological notes on *Conus* (Mollusca: Gastropoda) in the Trincomalee region of Ceylon. *Annals and Magazine of Natural History* (13) 2 (17): 309–320.

KOHN, A. J. 1960b. Spawning behaviour, egg masses and larval development in *Conus* from the Indian Ocean. *Bulletin of the Bingham Oceanographic Collection, Yale University* 17 (4): 1–51.

THIELE, J. 1910. Mollusca. B. Polyplacophora, Gastropoda marina, Bivalvia. In: SCHULTZE, L. *Zoologische und anthropologische Ergebnisse einer Forschungsreise im westlichen und zentralen Süd-Afrika ausgeführt in den Jahren 1903–1905* 4 (15). *Denkschriften der medizinisch-naturwissenschaftlichen Gesellschaft zu Jena* 16: 269–270.

(continued inside back cover)

ANNALS OF THE SOUTH AFRICAN MUSEUM
ANNALE VAN DIE SUID-AFRIKAANSE MUSEUM

Volume 98 Band
July 1988 Julie
Part 5 Deel



A REDESCRIPTION OF *AFROCHILTONIA*
CAPENSIS (K. H. BARNARD, 1916) WITH A
REVIEW OF THE GENERA OF THE FAMILY
CEINIDAE (CRUSTACEA, AMPHIPODA)

By
WOLFGANG ZEIDLER

Cape Town

Kaapstad

The ANNALS OF THE SOUTH AFRICAN MUSEUM

are issued in parts at irregular intervals as material
becomes available

Obtainable from the South African Museum, P.O. Box 61, Cape Town 8000

Die ANNALE VAN DIE SUID-AFRIKAANSE MUSEUM

word uitgegee in dele op ongereelde tye na gelang van die
beskikbaarheid van stof

Verkrygbaar van die Suid-Afrikaanse Museum, Posbus 61, Kaapstad 8000

OUT OF PRINT/UIT DRUK

1, 2(1-3, 5-8), 3(1-2, 4-5, 8, t.-p.i.), 5(1-3, 5, 7-9),
6(1, t.-p.i.), 7(1-4), 8, 9(1-2, 7), 10(1-3),
11(1-2, 5, 7, t.-p.i.), 14(1-2), 15(4-5), 24(2), 27, 31(1-3), 32(5), 33, 36(2), 45(1)

Copyright enquiries to the South African Museum

Kopieregnavrae aan die Suid-Afrikaanse Museum

ISBN 0 86813 096 6

Printed in South Africa by
The Rustica Press, Pty., Ltd.,
Court Road, Wynberg, Cape

In Suid-Afrika gedruk deur
Die Rustica-pers, Edms., Bpk.,
Courtweg, Wynberg, Kaap

A REDESCRIPTION OF *AFROCHILTONIA CAPENSIS*
(K. H. BARNARD, 1916)
WITH A REVIEW OF THE GENERA OF THE FAMILY CEINIDAE
(CRUSTACEA, AMPHIPODA)

By

WOLFGANG ZEIDLER

South Australian Museum, North Terrace, Adelaide, Australia

(With 4 figures)

[MS accepted 2 February 1988]

ABSTRACT

Afrochiltonia capensis (Barnard, 1916) is redescribed and figured, including the first description of the male, and a lectotype is selected. In the light of this redescription it is concluded that *A. capensis* is sufficiently different from currently recognized Australian congeners to warrant generic recognition. The genera of Ceinidae are reviewed briefly and *Austrochiltonia* Hurley, 1959, is resurrected for the Australian species with *Afrochiltonia* K. H. Barnard, 1955, being restricted to *A. capensis* from South Africa. A key to the genera of Ceinidae is provided.

CONTENTS

	PAGE
Introduction	105
Materials and methods	106
Systematics	107
Review of the family Ceinidae	114
Key to the genera of the family Ceinidae	118
Acknowledgements	118
References	118
Abbreviations	119

INTRODUCTION

Afrochiltonia capensis (Barnard, 1916) has never been adequately figured. The male was not correctly recognized until Griffiths (1976*b*) and has never been described, although R  he (1914), referring to six young females (as *Chiltonia subtenuis*), illustrated the second gnathopod of what is clearly a male.

Earlier confusion of non-ovigerous females with males led to a misleading generic diagnosis, which has never been resolved satisfactorily. Barnard (1916) suggested widening the diagnosis of *Chiltonia* to accommodate what he thought was a species with gnathopod one and two alike in both sexes. Later Hurley (1954) noted an unusual modification of the first pleopod in the male of New Zealand species of *Chiltonia*, where the inner ramus forms 1–3 dorsally directed whip-like lashes. This feature was thought to be of generic significance and led to

the erection of *Afrochiltonia* Barnard, 1955, for the South African species and *Austrochiltonia* Hurley, 1959, for the Australian species, with the New Zealand species remaining in *Chiltonia* Stebbing, 1899. Thus *Afrochiltonia* was distinguished from *Austrochiltonia* by the condition of the male second gnathopod. There the matter remained until Griffiths (1976*b*) discovered some males with the characteristically enlarged second gnathopod. He consequently synonymized *Afrochiltonia* with *Austrochiltonia* and furthermore *Afrochiltonia capensis* with *Austrochiltonia subtenuis*. However, *Afrochiltonia* has priority over *Austrochiltonia*, as noted and corrected by Barnard & Karaman (1982). Despite all of these systematic changes the South African species remained poorly known and the male was still not described or figured.

Barnard's (1916) description is considered inadequate by today's standards and is insufficient for comparisons with similar fauna in New Zealand and Australia. Because 'Chiltonias' are very common and widespread in the freshwaters of southern Australia, with several undescribed species, it is important to establish clearly the systematic status of the South African species.

It is the purpose of this paper to redescribe *Afrochiltonia capensis* (Barnard, 1916), including the first description of the male, and to clarify the status of the genera of Ceinidae, particularly *Afrochiltonia* and *Austrochiltonia*.

MATERIALS AND METHODS

Barnard's (1916) type specimens of *Afrochiltonia capensis* were borrowed from the South African Museum and examined in detail. As no other material of *A. capensis* was available from the South African Museum, additional specimens were collected at my request, from Milnerton Lagoon (see 'Material examined') by Dr C. L. Griffiths. These specimens are deposited in the South Australian Museum, except for the male described herein, which has been transferred to the South African Museum. Specimens of *Austrochiltonia* in the collections of the South Australian Museum were also examined for comparison.

Specimen length was measured along a lateral parabolic line drawn from the anterior extremity of the head through the middle of the body to the posterior limit of the telson. Barnard's type material was not used for size comparisons between males and females as his sample may have been biased towards larger specimens.

The thoracic limbs are referred to as gnathopod 1 and 2 followed by pereopods 3–7 to avoid confusion. Size comparisons of gnathopods exclude the coxa and dactyl and of the pereopods the coxa, with articles being measured down the middle.

Unless indicated otherwise dissected appendages were taken from the left hand side of the animal. The mouthparts of the lectotype and the mouthparts and appendages from the male, described herein, are mounted in poly-vinyl lactophenol on microscope slides. All other appendages, remains of dissected specimens, and other specimens are preserved in 75 per cent alcohol.

SYSTEMATICS

Family **Ceinidae** J. L. Barnard, 1972*Afrochiltonia capensis* (Barnard, 1916)

Figs 1–4

Chiltonia subtenuis Rühe, 1914: 35, figs 13, 14a–c [non Sayce, 1902].*Chiltonia capensis* Barnard, 1916: 224, pl. 27 (figs 38–40).*Afrochiltonia capensis* Barnard, 1955: 93. Griffiths, 1974a: 253; 1974b: 327; 1975: 168; 1976a: 75, fig. 47.*Austrochiltonia subtenuis* Griffiths, 1976b: 30.*Type locality*

Salt River, Cape Town, South Africa, by present designation of lectotype.

*Material examined**Type material.* K. H. Barnard's syntypes consist of two lots.

SAM–A2885 labelled 'Type Specms' from Salt River, Cape Town, collected by Dr W. F. Purcell, October 1898, consisting of 8 females in alcohol, one of which has been selected as lectotype; the remainder have been designated paralectotypes and have been transferred to SAM–A39685.

SAM–A2886 from Milnerton near Cape Town, collected by K. H. Barnard, 25 October 1913, consisting of 20 females (some damaged) in alcohol and a microscope slide of the appendages of at least three specimens. All of this material has been designated paralectotypes. The slide material, although labelled 'Type', could not be used as the lectotype as it did not consist of a single specimen and the mountant had become crazed with age, thus obscuring the finer detail of the mounted appendages.

Other material. South Australian Museum No. C4165: 30 females and 6 males from Milnerton Lagoon, Cape Town, collected by C. L. Griffiths, 20 January 1987. SAM–A39686: male (the specimen described and illustrated herein), collection data as for C4165 but transferred to the South African Museum. South Australian Museum No. C4166: 40 females and 69 males from mouth of Milnerton Lagoon, near Cape Town, collected by C. L. Griffiths, June 1987.

*Description**Female*

Lectotype, 3,7 mm, non-ovigerous, SAM–A2885. Coxal gills present from G2 to P6. Oostegites, dorsally folded, present from G2 to P5.

Head as long as deep, length equivalent to first 1,5 pereonites; eyes black (in alcohol), ovato-circulate with some ommatidia diffuse dorsally.

Antenna 1 short, about twice head length or equivalent to 0,2 times body length; article 1 of peduncle almost twice as long as wide and 1,5 times length of article 2; article 3 slightly shorter than 2; flagellum slightly longer than peduncle, of six articles with a ventral aesthetasc at the base of each of the last two articles.




Fig. 1. *Afrochiltonia capensis* (Barnard). A. Lectotype female, 3,7 mm (SAM-A2885) (gills, oostegites and pleopods not shown). Scale bar = 0,5 mm. B. Oostegites from paralectotype specimen (SAM-A39685), shown in order from G2-P5. Scale bar = 0,2 mm. Other appendages from lectotype. Scale bar = 0,2 mm.




Fig. 2. *Afrochiltonia capensis* (Barnard). Lectotype female, 3.7 mm, gnathopods and mouthparts. Scale bars = 0.2 mm and 0.1 mm, respectively.

Antenna 2 little more than 0,75 length A1, gland cone pressed firmly against head—not visible laterally; article 1 of peduncle twice as wide as long; article 2 as wide as long, twice length article 1 and slightly less than half length of article 3; flagellum only slightly longer than peduncle, of six articles.

Upper lip as wide as long, apically rounded, bearing numerous short setae distally.

Lower lip without inner lobes; outer lobes subovate with setose distal and inner margins.

Mandibles without palp: left with incisor of seven teeth, lacina mobilis of five teeth, spine row of three feathered spines and triturative molar with one long feathered seta; right with incisor of five teeth, lacina mobilis of three teeth, spine row of three feathered spines and molar like left.

Maxilla 1: outer plate without palp but notched at palp's normal position, with eight comb-like spines apically; inner plate very narrow with only two feathered spines apically.

Maxilla 2: outer plate slightly longer than inner, about 0,75 times as wide; both apically setose; inner plate with one large seta on inner margin at end of setal row.

Maxilliped: inner plate reaching extremity of article 1 of palp, rectangular, about three times as long as broad, with three stout spines apically, the inner one very small; outer plate reaching 0,75 along article 2 of palp, ovate, about as wide as inner plate bearing several setae apically and along inner margin; palp article 1 with oblique distal margin, length outer margin about 2,5 times inner; palp article 2 slightly broader than long and slightly shorter than outer margin of article 1, bearing a few setae on inner distal corner and distal half of inner margin; palp article 3 about as long as broad and as long as article 2, with sparse long setae on distal and inner margins; palp article 4 small, conical, slightly longer than wide, about half length of article 3; dactylus sharp, as long as article 4.

Gnathopod 1: coxa length about 1,5 times width, longer than article 2 with antero-dorsal corner slightly produced; article 5 length twice maximum width, postero-distal lobe not produced, with row of six stout pectinate spines; article 6 as long as article 5, about twice as long as wide, postero-distal corner rounded, distal face with spine on either side of dactyl and two long medial setae, antero-distal corner with four long setae, posterior margin with small spine near postero-distal corner; dactyl as long as width of palm and fitting neatly against palm.

Gnathopod 2: similar to G1 but 1,25 times as long; coxal gill sac-like, more than twice as long as wide, as long as article 2; coxa slightly longer than wide, as long as article 2; article 5 with slightly produced postero-distal lobe with row of five stout pectinate spines; article 6 without small spine on posterior margin, otherwise as in G1.

Pereopod 3: length 1,35 times G2; coxal gill like that of G2; coxa like that of G2 but slightly larger, slightly longer than article 2; article 4 broad, about

0,7 times length; article 5 like 4 but not as broad and only 0,75 times as long; article 6 length 1,6 times article 5, about three times as long as wide; dactyl length 0,4 times article 6; all articles sparsely setose as illustrated.

Pereopod 4 identical to P3 except for coxa. Coxa with shallow antero-dorsal excavation, maximum width 1,5 times length, slightly longer than article 2, posterior margin oblique so that width at distal margin is only about half maximum width.

Pereopod 5: smallest pereopod, length about 0,9 times P4; coxal gill similar to P4 but a little wider; coxa width almost twice width article 2, length of anterior lobe about half maximum width coxa, length of posterior lobe about 0,7 maximum width coxa or as long as article 2; article 2 slightly longer than wide with typical expanded posterior margin and postero-distal lobe overlapping and almost reaching to distal margin of article 3; article 4 length 1,2 times width, with postero-distal corner produced; article 5 length about 1,2 times article 4 and of similar shape except postero-distal corner is not as produced; article 6 length 1,6 times article 5, about three times as long as wide; dactyl length 0,4 times article 6; all articles sparsely spinose as illustrated.

Pereopod 6 longest pereopod, length 1,25 times P5; like P5 except articles 3–6 somewhat longer in proportion to their width; coxa as wide as article 2, anterior lobe small, length about half width coxa, posterior lobe as long as coxa width.

Pereopod 7 a little shorter than P6; coxa semi-circular, slightly wider than long; article 2 as wide as long, postero-distal lobe extending beyond article 3, posterior margin slightly serrate and minutely spined with acute proximal corner; otherwise similar to P6.

Pleonal epimera with very small postero-ventral tooth.

Uropod 1 longer than U2; rami subequal, about 0,8 times as long as peduncle, outer ramus with two large and two small spines at tip, inner ramus with two large and three small spines at tip; peduncle with large spine on inner and outer-distal corner and two more on dorsal outer margin.

Uropod 2: rami subequal, a little more than 0,8 times as long as peduncle; peduncle and rami with spines as in U1 except outer ramus also has a spine on middle of inner margin (only on right in lectotype).

Uropod 3 one-articulate, half length of telson, conical in shape with one long outer and one short inner seta at tip.

Telson entire, hemispherical, slightly wider than long.

Oostegites from paralectotype (SAM-A39685), ovigerous, most eggs released. All with curled margins and numerous small hooks forming a tight marsupium. First is pentagonal with long distal and posterior margins, almost as wide as deep; second is rectangular, length 1,8 times width; third is also rectangular, length a little more than 1,8 times width and is longest oostegite, about 0,6 times length of P4; fourth is sub-rectangular with oblique distal margin and excavate postero-distal corner, maximum length 1,3 times maximum width.

Male

Hypotype 2,5 mm (SAM-A39686), generally like female but differs as follows.

Coxal gills relatively smaller.

Antenna 1 flagellum a little shorter than peduncle, consisting of only five articles.

Antenna 2 only slightly shorter than A1.

Mandibles: only right molar with long feathered seta.

Gnathopods and pereopods with articles not so stout.

Gnathopod 1: coxa narrower distally, about 0,7 times dorsal width, without antero-dorsal corner produced; article 6 slightly longer than article 5 with two spines on distal face in addition to those on either side of the dactyl.

Gnathopod 2 with enlarged article 6, unlike G1; length about 1,4 times G1; coxa length 1,2 times width, only 0,9 times length article 2; article 4 with right-angled bend; article 5 small, without pectinate spines; article 6 a little longer than article 2, maximum length 1,6 times maximum width, postero-proximal corner forming distinct lobe for almost 0,4 length article, palm oblique with several small spines on either side of 'cutting edge' followed proximally by small groove for tip of dactyl; dactyl claw-like, length 0,8 times maximum length article 6.

Pereopod 3: length 1,1 times G2; coxa like that of G2 only slightly longer.

Pereopod 4: coxa maximum width a little more than length, only slightly more narrow distally—posterior corner of excavation not produced as in female.

Pereopod 5 as long as P4; coxa width 1,5 times width article 2, length anterior lobe slightly less than half maximum width coxa; length posterior lobe less than 0,6 times maximum width coxa or only 0,7 times length article 2.

Pereopod 6: coxa, length anterior lobe 0,3 times width coxa, length posterior lobe a little less than coxa width.

P7, U1–U3 and telson like female.

Pleopods all of normal structure (not modified as in *Chiltonia*).

Variations

Females ranged in size from 1,0 to 4,1 mm with a mean of $3,2 \pm 0,2$ mm ($\pm 95\%$ c.l.; $n = 70$). Males ranged in size from 1,5 to 2,6 mm with a mean of $2,2 \pm 0,06$ mm ($\pm 95\%$ c.l.; $n = 76$). Males are thus significantly smaller than females ($P < 0,01$). No specimens were as large as 4,5 mm as recorded by Barnard (1916) and one must assume this measurement included the antennae or to be an error.

All of the specimens examined varied little from the above descriptions. In some specimens the ommatidia of the eyes were more diffuse at the edges and almost confluent dorsally. Antenna 1 and 2 usually had a flagellum of six articles, increasing to seven in some females, or decreasing to five in some males but rarely in females. The antennae were almost equal in length in some specimens and varied from 0,2 to 0,25 times the body length. The condition of the mandible with a feathered seta on the molar usually occurred only on the right, rarely on




Fig. 3. *Afrochiltonia capensis* (Barnard). Lectotype female, 3,7 mm, pereopods 1-7.
Scale bar = 0,2 mm.

the left as in the lectotype. The rami of U1 were sometimes as short as 0.6 times peduncle, particularly in males. The inner ramus of U2 without a medial spine was rare. Oostegites of females varied considerably in size but were expanded and as illustrated in ovigerous specimens.

Remarks

Griffiths (1976b) synonymized *Afrochiltonia capensis* with *Austrochiltonia subtenuis* on the basis that the male G2 was enlarged and U3 was one-articulate. However, apart from a number of minor differences, *Afrochiltonia capensis* is clearly distinguished from *Austrochiltonia subtenuis* by: (1) the shape of coxa 4; (2) the relative lengths of the antennae; (3) the stout nature of the pereopods, particularly in the female; (4) the lack of marginal spines on the rami of U1 and U2; (5) the gland cone on A2 is not visible laterally; and (6) P7 is shorter than P6. The synonymy proposed by Griffiths (1976b) is thus considered invalid.

The earlier confusion on non-ovigerous females with males might be explained by the fact that the males are considerably smaller than the females and may have been overlooked when sorting samples or confused with juveniles of other common species (e.g. *Melita zeylanica*). There is also a likely seasonal variation in the number of males present in the population, as is evidenced by the two random samples collected in January and June 1987, which contained 6/36 and 69/109 males, respectively. It is therefore possible that, when Barnard's samples were collected in October, few males were present in the natural population and were thus not represented in his samples. Further evidence of a seasonal breeding cycle is provided by the fact that in January 15/30 females were ovigerous as compared to only 9/40 in June. However, a more detailed study of the life cycle of this species is required to determine breeding seasons and seasonal variations in the male/female ratio.

Now that the male has been described it is possible to re-evaluate the systematic position of this species with that of similar taxa in New Zealand and Australia. In particular, I recommend retaining the genus *Austrochiltonia* Hurley, 1959, for the Australian species and reserving *Afrochiltonia* Barnard, 1955, for the single South African species.

Distribution

Kosi Bay, Zululand, to Olifants River, western Cape, in brackish-estuarine environments (endemic).

REVIEW OF THE FAMILY CEINIDAE J. L. BARNARD, 1972

The family Ceinidae is divided into two subfamilies, the Ceininae—consisting of marine forms with cleft telsons—and the Chiltoniinae—consisting of freshwater-brackish forms with uncleft telsons (Barnard 1972b). It is generally distinguished from the Hyalidae (= Talitridae) by the form of U3 (e.g. Barnard 1972b), which is diagnosed by Barnard (1972a) as 'composed only of peduncle




Fig. 4. *Afrochiltonia capensis* (Barnard). Male, 2,5 mm (SAM-A39686), pereopods 1-7.
Scale bar = 0,2 mm.